



CLAIMS:

1. An improved for the preparation of (S)- atenolol (1), comprising the steps of:


a) reacting a phenol of formula 2:




with an (R)-epichlorohydrin of formula (3) :



in presence of an alkali metal hydroxide and a quaternary ammonium salt as phase transfer catalyst in an aqueous solution at a temperature of -10°C to 0°C to obtain optically active intermediate glycidyl ether of formula 4:



b) reacting the optically active intermediate glycidyl ether (4) with isopropylamine at 10°C to 40°C to obtain (S)-atenolol of formula 1:



in high optical purity of >99 ee.

2. A process as claimed in claim 1 wherein the alkali metal hydroxide is selected from sodium hydroxide or potassium hydroxide.
3. A process as claimed in claim 1 wherein the amount of alkali metal hydroxide is 1 to 1.5 moles to 1 mole of the phenol (2).
4. A process as claimed in claim 1 wherein the amount of (R)-epichlorohydrin is 1 to 3 moles to 1 mole of the phenol (2).
5. A process as claimed in claim 1 wherein the quaternary ammonium salt has the formula $R^1R^2R^3R^4N^+X^-$

Wherein R^1 , R^2 , R^3 and R^4 are same or different and are alkyl groups having 1 to 16 carbon atoms selected from methyl, ethyl, propyl butyl, phenyl or benzyl, X is a group selected from chlorine, bromine, iodine, hydrogen sulphate or hydroxyl group.

6. A process as claimed in claim 1 wherein the amount of quaternary ammonium salt is 0.001 to 2% by weight of phenol (2).
7. A process as claimed in claim 1 further comprising formation of chlorohydrine (5) as side product.
8. A process as claimed in claim 1 further comprising reacting chlorohydrine (5) with isopropylamine at 10 to 40°C to obtain S-atenolol.